

University of Groningen

Ancient goat genomes reveal mosaic domestication in the Fertile Crescent

Daly, Kevin G.; Delser, Pierpaolo Maisano; Mullin, Victoria E.; Scheu, Amelie; Mattiangeli, Valeria; Teasdale, Matthew D.; Hare, Andrew J.; Burger, Joachim; Verdugo, Marta Pereira; Collins, Matthew J.

Published in:
Science Magazine

DOI:
[10.1126/science.aas9411](https://doi.org/10.1126/science.aas9411)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Daly, K. G., Delser, P. M., Mullin, V. E., Scheu, A., Mattiangeli, V., Teasdale, M. D., Hare, A. J., Burger, J., Verdugo, M. P., Collins, M. J., Kehati, R., Erek, C. M., Bar-Oz, G., Pompanon, F., Cumer, T., Cakirlar, C., Mohaseb, A. F., Decruyenaere, D., Davoudi, H., ... Bradley, D. G. (2018). Ancient goat genomes reveal mosaic domestication in the Fertile Crescent. *Science Magazine*, 361(6397), 85-87.
<https://doi.org/10.1126/science.aas9411>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

DOMESTICATION

Ancient goat genomes reveal mosaic domestication in the Fertile Crescent

Kevin G. Daly^{1*}, Pierpaolo Maisano Delser^{1,2*}, Victoria E. Mullin^{1,3}, Amelie Schen^{1,4}, Valeria Mattiangeli¹, Matthew D. Teasdale^{1,5}, Andrew J. Hare¹, Joachim Burger⁴, Marta Pereira Verdugo¹, Matthew J. Collins^{5,6}, Ron Kehati⁷, Cevdet Merih Ereğ⁸, Guy Bar-Oz⁹, François Pompanon¹⁰, Tristan Cumer¹⁰, Canan Çakırlar¹¹, Azadeh Fatemeh Mohaseb^{12,13}, Delphine Decruyenaere¹², Hossein Davoudi^{14,15}, Özlem Çevik¹⁶, Gary Rollefson¹⁷, Jean-Denis Vigne¹², Roya Khazaeli¹³, Homa Fathi¹³, Sanaz Beizae Doost¹³, Roghayeh Rahimi Sorkhani¹⁸, Ali Akbar Vahdati¹⁹, Eberhard W. Sauer²⁰, Hossein Azizi Kharanaghi²¹, Sepideh Maziar²², Boris Gasparian²³, Ron Pinhasi²⁴, Louise Martin²⁵, David Orton⁵, Benjamin S. Arbuckle²⁶, Norbert Benecke²⁷, Andrea Manica², Liora Kolska Horwitz⁷, Marjan Mashkour^{12,13,15}, Daniel G. Bradley^{1†}

Current genetic data are equivocal as to whether goat domestication occurred multiple times or was a singular process. We generated genomic data from 83 ancient goats (51 with genome-wide coverage) from Paleolithic to Medieval contexts throughout the Near East. Our findings demonstrate that multiple divergent ancient wild goat sources were domesticated in a dispersed process that resulted in genetically and geographically distinct Neolithic goat populations, echoing contemporaneous human divergence across the region. These early goat populations contributed differently to modern goats in Asia, Africa, and Europe. We also detect early selection for pigmentation, stature, reproduction, milking, and response to dietary change, providing 8000-year-old evidence for human agency in molding genome variation within a partner species.

The Fertile Crescent of Southwest Asia and adjacent areas were the location of transformative prehistoric innovations including the domestication of sheep, goats, cattle, and pigs (1–3). Archaeological evidence suggests local development of wild goat (bezoar) management strategies in different regions in the mid- to late 11th millennium before the present (BP) with domestic phenotypes emerging in the 10th millennium, first in the Anatolian region

(4–6). A key question is whether these early patterns of exploitation are consistent with a geographically focused singular domestication process or whether domestic goats were recruited from separate populations, with parallel genetic consequences. Genetic evidence is inconclusive (7, 8).

We generated ancient *Capra* genome data from Neolithic sites from western (Anatolia and the Balkans), eastern (Iran and Turkmenistan), and southern (Jordan and Israel) regions around the

Fertile Crescent (tables S1 to S3). To maximize yields, we sampled mainly petrous bones; 51 such samples produced nuclear genome coverage ranging from 0.01× to 14.89× (median 1.05×) (tables S4 and S5). We enriched for mitochondrial DNA (mtDNA) in poorly preserved samples and obtained a total of 83 whole mitochondrial genomes (median 70.95×) (table S6 and figs. S1 and S2) (9).

The majority of our ancient domestic mitochondrial sequences fall within modern haplogroups A, B, C, D, and G (Fig. 1A, figs. S3 to S6, and tables S7 to S9). The Paleolithic wild goat samples fall exclusively in more divergent clades T [similar to the related wild caprid, the West Caucasian tur (*Capra caucasica*)] and F [previously reported in bezoar and a small number of Sicilian goats (10)]. Here, we found F in a >47,000 BP bezoar from Hovk-1 cave, Armenia; in a pre-domestic goat from Direkli Cave, Turkey; and in Levantine goats at 'Ain Ghazal, an early Neolithic village in Jordan, and Abu Ghosh, Israel.

A geographic plot of Neolithic samples illustrates that early domestic goat haplogroups are highly structured (Fig. 1B), with disjunct distributions in the western, eastern, and southern (Levantine) regions of the Near East (tables S10 and S11). In this early farming period, partitioning is significant; analysis of molecular variance (9) estimates that 81% of the mtDNA diversity stems from differences between the three regions ($P = 0.028$, permutation test) (tables S12 and S13). When we use an approximate Bayesian computation (ABC) framework on this mtDNA variation to investigate demographic history, a model suggesting a pre-domestic branching of the divergent Levant population (38,500 to 195,200 BP) is favored. This suggests multiple wild origins of Neolithic goat herds (tables S14 to S19) (9). In the later post-Neolithic samples,




Fig. 1. Maximum likelihood phylogeny and geographical distributions of ancient mtDNA haplogroups. (A) A phylogeny placing ancient whole mtDNA sequences in the context of known haplogroups. Symbols denoting individuals are colored by clade membership; shape indicates archaeological period (see key). Unlabeled nodes are modern bezoar and outgroup sequence (Nubian ibex) added for reference. We define haplogroup T as the sister branch to the West Caucasian tur (9). (B and C) Geographical distributions of haplogroups show early highly structured diversity in the Neolithic period (B) followed by collapse of structure in succeeding periods (C). We delineate the tiled maps at 7250 to 6950 BP, a period bracketing both our earliest Chalcolithic sequence (24, Mianroud) and latest Neolithic (6, Aşağı Pınar). Numbered archaeological sites also include Direkli Cave (8), Abu Ghosh (9), 'Ain Ghazal (10), and Hovk-1 Cave (11) (table S1) (9).

this partitioning collapses to zero (Fig. 1C) and the ubiquitous modern haplogroup, A, becomes widespread.

Analyses of genome-wide variation also argue against a single common origin. Neolithic samples from the west, east, and Levant each cluster separately in principal components analysis (PCA; Fig. 2) and in phylogenetic reconstruction (figs. S7 to S10). *D* statistics show that these clusters have significantly different levels of allele sharing with two regional samples of pre-domestic wild goats: a ~13,000 BP population from Direkli cave (Southeast Anatolia) and a >47,000 BP bezoar from Hovk-1 cave (Armenia) (Fig. 3A) (9). These differences are consistent with qpGraph estimation of relationships (Fig. 3B, fig. S11, and table S20) where a primary ancestral divide between western and eastern genomes occurred more than 47,000 BP. The latter clade gave rise to the eastern Neolithic population. However, the western and Levant Neolithic goat derive ~50% and ~70% of their ancestry from a divergent source in the western clade that had affinity to the Anatolian wild population, in line with f_4 ratios and Treemix graphs (fig. S12 and table S21). These different proportions infer substantial local recruitment from different wild populations into early herds in regions proximal to each of the different vertices of the Fertile Crescent. ABC modeling of autosomal variation also rejects a single domestication origin scenario (figs. S13 to S15 and tables S11 and S22 to S25) (9).

Thus, our data favor a process of Near Eastern animal domestication that is dispersed in space and time, rather than radiating from a central core (3, 11). This resonates with archaeozoological evidence for disparate early management strategies from early Anatolian, Iranian, and Levantine Neolithic sites (12, 13). Interestingly, our finding of divergent goat genomes within the Neolithic echoes genetic investigation of early farmers. Northwestern Anatolian and Iranian human Neolithic genomes are also divergent (14–16), which suggests the sharing of techniques rather than large-scale migrations of populations across Southwest Asia in the period of early domestication. Several crop plants also show evidence of parallel domestication processes in the region (17).

PCA affinity (Fig. 2), supported by qpGraph and outgroup f_3 analyses, suggests that modern European goats derive from a source close to the western Neolithic; Far Eastern goats derive from




Fig. 2. Principal components analysis of ancient and modern goat genomes. Ancient goats cluster in three vertices: eastern (Iran, Uzbekistan, Turkmenistan, Georgia), western (Balkans, Anatolia), and southern or Levantine (Jordan, Israel) margins of the Near East. Modern European, Asian and, interestingly, African goats follow this pattern, but Bronze Age Anatolian (red arrow) and Chalcolithic/Bronze Age Israeli (yellow arrow) samples show shifts relative to earlier genomes from those regions, suggesting post-Neolithic admixture within the primary regions.

early eastern Neolithic domesticates; and African goats have a contribution from the Levant, but in this case with considerable admixture from the other sources (figs. S11, S16, and S17 and tables S26 and S27). The latter may be in part a result of admixture that is discernible in the same analyses extended to ancient genomes within the Fertile Crescent after the Neolithic (figs. S18 and S19 and tables S20, S27, and S31) when the spread of metallurgy and other developments likely resulted in an expansion of inter-regional trade networks and livestock movement.

Animal domestication likely involved adaptive pressures due to infection, changes in diet, translocation beyond natural habitat, and human selection (18). We thus took an outlier approach to identify loci that underwent selective sweeps in either six eastern Neolithic genomes or four

western genome samples (minimum coverage 2×). We compared each population to 16 modern bezoar genomes (19) and identified 18 windows with both high divergence (highest 0.1% *Fst* values) and reduced diversity in Neolithic goats (lowest 5% θ ratio: Neolithic/wild; tables S28, S29, and S32).

The pigmentation loci, *KIT* and *KITLG*, are the only shared signals in both Neolithic populations. Both are common signals in modern livestock analyses (19, 20). We thus examined *Fst* values for previously reported coloration genes and identified *ASIP* and *MITF* as also showing high values (Fig. 4, A and B, fig. S20, and table S30). Whereas modern breeds are defined in part by color pattern, the driver of the ~8000-year-old selection observed in the Neolithic for pigmentation may be less obvious. *KIT* is involved in the piebald trait in mammals (21) and may have been favored as a means of distinguishing individuals and maintaining ownership within shared herds as well as for aesthetic value. Pigmentation change has also been proposed as a pleiotropic effect of selection for tameness (22). Intriguingly, selective sweeps around the *KIT* locus were clearly independent in the eastern and western Neolithic goat sampled genomes, as the resulting locus genotypes are distinct and contribute differently to modern eastern and western populations (Fig. 4C).

Trait mapping in cattle, the most studied ungulate, offers interpretation of three other caprine signals identified here. *SIRT1* (identified in the western Neolithic) has variants affecting stature (23), and a reduction in size is a widespread signal of early domestication. *EPGN* (eastern Neolithic) is linked to calving interval; increase in reproductive frequency is another general feature of domestication. *STAT1* (eastern Neolithic) is involved in mammary gland development and has been linked to milk production (24). The second most extreme eastern signal maps to a homolog of human *CYP2C19*, which (like other cytochrome P450 products) contributes to metabolism of xenobiotics including enniatin B, a toxic product of fungal strains that contaminate cereals and grains. This selection signal has been hypothesized as a response to early agriculture in humans (25). Early recycling of agricultural by-products as animal fodder has been suggested as a motivation for the origins of husbandry (3), and fungal toxins may have been a challenge to early domestic goats as well as their agriculturist owners.

Our results imply a domestication process carried out by humans in dispersed, divergent,

¹Smurfit Institute of Genetics, Trinity College Dublin, Dublin 2, Ireland. ²Department of Zoology, University of Cambridge, Cambridge CB2 3EJ, UK. ³Department of Earth Sciences, Natural History Museum, London SW7 5BD, UK. ⁴Palaeogenetics Group, Institute of Organismic and Molecular Evolution (iOME), Johannes Gutenberg University Mainz, 55099 Mainz, Germany. ⁵BioArCh, University of York, York YO10 5DD, UK. ⁶Museum of Natural History, University of Copenhagen, Copenhagen, Denmark. ⁷National Natural History Collections, Faculty of Life Sciences, The Hebrew University, Jerusalem, Israel. ⁸Gazi University, Ankara 06500, Turkey. ⁹Zinman Institute of Archaeology, University of Haifa, Mount Carmel, Haifa, Israel. ¹⁰Université Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LECA, F-38000 Grenoble, France. ¹¹Groningen Institute of Archaeology, Groningen University, Groningen, Netherlands. ¹²Archéozoologie, Archéobotanique (UMR 7209), CNRS, MNHN, UPMC, Sorbonne Universités, Paris, France. ¹³Archaeozoology section, Archaeometry Laboratory, University of Tehran, Tehran, Iran. ¹⁴Department of Archaeology, Faculty of Humanities, Tarbiat Modares University, Tehran, Iran. ¹⁵Osteology Department, National Museum of Iran, Tehran, Iran. ¹⁶Trakya Üniversitesi, Edebiyat Fakültesi, Arkeoloji Bölümü, Edirne, Turkey. ¹⁷Department of Anthropology, Whitman College, Walla Walla, WA 99362, USA. ¹⁸Faculty of Cultural Heritage, Handicrafts and Tourism, University of Mazandaran, Noshahr, Iran. ¹⁹Provincial Office of the Iranian Center for Cultural Heritage, Handicrafts and Tourism Organisation, North Khorassan, Bojnord, Iran. ²⁰School of History, Classics and Archaeology, University of Edinburgh, William Robertson Wing, Old Medical School, Edinburgh EH8 9AG, UK. ²¹Prehistory Department, National Museum of Iran, Tehran, Iran. ²²Institut für Archäologische Wissenschaften, Goethe Universität, Frankfurt am Main, Germany. ²³Institute of Archaeology and Ethnology, National Academy of Sciences of the Republic of Armenia, Yerevan 0025, Republic of Armenia. ²⁴Department of Anthropology, University of Vienna, 1090 Vienna, Austria. ²⁵Institute of Archeology, University College London, London, UK. ²⁶Department of Anthropology, University of North Carolina, Chapel Hill, NC, USA. ²⁷Department of Natural Sciences, German Archaeological Institute, 14195 Berlin, Germany.

*These authors contributed equally to this work.
†Corresponding author. Email: dbradley@tcd.ie




Fig. 3. *D* statistics and admixture graph of ancient and modern goats. (A) In the test $X(Y, Z)$, positive or negative D values indicate a greater number of derived alleles between X and Z or X and Y , respectively; yak is used as an outgroup. D values for each test are presented with error bars of 3 SE; a nonsignificant test is shown in gray. These show that regional pre-domestic wild goats relate asymmetrically to Neolithic domestic populations, ruling out a singular origin. (B) Admixture graph reconstructing the population history of pre-Neolithic and Neolithic goats. Relative inputs from divergent sources into early domestic herds are represented by gray dashed arrows (drawn from fig. S11F) (9).




Fig. 4. *Fst* distributions between modern bezoar and Neolithic western and eastern populations, and a heat map of allele sharing between modern and domestic goats at the *KIT* locus. (A and B) The highest F_{st} values for 50-kb windows overlapping seven pigmentation loci showing evidence of selection in modern goat, sheep, or cattle studies are indicated for western (A) and eastern (B) populations (tables S30 and S32). (C) The pigmentation locus, *KIT*, shows evidence of selection in both western and eastern Neolithic samples, but allele sharing distances (illustrated as a heat map) suggest that selection acted on divergent standing variation in parallel but separate processes. Five of the seven ancient western samples are from Neolithic contexts and cluster with modern western haplogroups. The two remaining western ancients (red) falling in the eastern cluster (mainly blue) are Bronze Age Anatolian samples with indications of secondary admixture (Fig. 2).

but communicating communities across the Fertile Crescent who selected animals in early millennia, including for pigmentation, the most visible of domestic traits.

REFERENCES AND NOTES

- J. Peters, A. von den Driesch, D. Helmer, in *The First Steps of Animal Domestication: New Archaeological Approaches*, J. D. Vigne, J. Peters, D. Helmer, Eds. (Oxbow, 2005), pp. 96–123.
- M. A. Zeder, *J. Anthropol. Res.* **68**, 161–190 (2012).
- J.-D. Vigne, L. Gourichon, D. Helmer, L. Martin, J. Peters, in *Quaternary in the Levant*, Y. Enzel, O. Bar Yosef, Eds. (Cambridge Univ. Press, 2017), pp. 753–760.
- M. A. Zeder, B. Hesse, *Science* **287**, 2254–2257 (2000).
- D. Helmer, L. Gourichon, in *Archaeozoology of the Near East*, vol. 9, M. Mashkour, M. Beech, Eds. (Oxbow, 2017), pp. 23–40.
- B. Moradi et al., in *The Neolithic of the Iranian Plateau*, K. Roustaei, M. Mashkour, Eds. (Ex Oriente, 2016), pp. 1–14.
- S. Naderi et al., *Proc. Natl. Acad. Sci. U.S.A.* **105**, 17659–17664 (2008).
- P. Gerbault et al., in *Population Dynamics in Prehistory and Early History: New Approaches Using Stable Isotopes and Genetics*, E. Kaiser, J. Burger, W. Schier, Eds. (De Gruyter, 2012), pp. 17–30.
- See supplementary materials.
- M. T. Sardina et al., *Anim. Genet.* **37**, 376–378 (2006).
- M. A. Zeder, *Curr. Anthropol.* **52**, S221–S235 (2011).
- L. K. Horwitz et al., *Paléorient* **25**, 63–80 (1999).
- B. S. Arbuckle, L. Atici, *Levant* **45**, 219–235 (2013).
- F. Broushaki et al., *Science* **353**, 499–503 (2016).
- I. Lazaridis et al., *Nature* **536**, 419–424 (2016).
- M. Gallego-Llorente et al., *Sci. Rep.* **6**, 31326 (2016).
- D. Q. Fuller, G. Willcox, R. G. Allaby, *World Archaeol.* **43**, 628–652 (2011).
- M. A. Zeder, *Interface Focus* **7**, 20160133 (2017).
- F. J. Alberto et al., *Nat. Commun.* **9**, 813 (2018).
- J. W. Kijas et al., *PLOS Biol.* **10**, e1001258 (2012).
- N. Reinsch et al., *J. Hered.* **90**, 629–634 (1999).
- L. Trut, I. Oskina, A. Kharlamova, *BioEssays* **31**, 349–360 (2009).
- M. Li et al., *Mol. Cell. Probes* **27**, 215–220 (2013).
- O. Cobanoglu, I. Zaitoun, Y. M. Chang, G. E. Shook, H. Khatib, *J. Dairy Sci.* **89**, 4433–4437 (2006).
- R. E. Janha et al., *BMC Evol. Biol.* **14**, 71 (2014).

ACKNOWLEDGMENTS

We thank L. Cassidy, E. Jones, L. Frantz, G. Larson, and M. Zeder for critical reading of the text; the Israel Antiquities Authority for permitting sampling of the Israeli sites (under permit); excavators, archaeozoologists, and museums who permitted sampling from their excavations and collections without which this project would not have been possible, including T. Levy, C. Grigson, A. Maeir, S. Gitin, P. de Miroschedji, S. Davis, and A. Ben-Tor; the Iranian Cultural Heritage Handicraft and Tourism organization and the National Museum of Iran (NMI) and J. Nokandeh, director, and F. Biglari, cultural deputy; H. Laleh and A. Aliyari, Directors of the Archaeometry Laboratory of the University of Tehran. The ATM Project of MNHN supported sampling of several sites as well as the LIA HAOMA CNRS project. We are grateful for assistance from J. Vuković, J. Bulatović, I. Stojanović, H. Greenfield, Wiltshire Museum, L. Brown, and Trinseq. The authors wish to acknowledge the DJEI/DES/SFI/HEA Irish Centre for High-End Computing (ICHEC) for the provision of computational facilities and support. **Funding:** Supported by ERC Investigator grant 295729-CodeX. Additional support from Science Foundation Ireland Award 12/ERC/B2227. P.M.D. was supported by the HERA Joint Research Programme “Uses of the Past” (CitiGen); and the European Union’s Horizon 2020 research and innovation program under grant agreement no. 649307. A.M. was supported by ERC Consolidator grant 647787-LocalAdaptation. M.D.T. was supported by the Marie Skłodowska-Curie Individual Fellowship SCRIBE H2020-MSCA-IF-2016 747424. Author contributions: D.G.B. conceived of the project and designed the research, with input from J.B. and M.C.; C.C., R.P., L.M., D.O., B.S.A., N.B., L.K.H., M.M., R.Ke., C.M.E., G.B.O., F.P., T.C., J.D.V., A.F.M., D.D., H.D., Ö.C., R.Kh., H.F., S.B., R.R.S., A.A.V., E.W.S., H.A.K., and S.M. provided samples and data; K.G.D., V.E.M., V.M., A.S., A.J.H., and M.D.T. performed genomics laboratory work; P.M.D. performed ABC analyses, with input from A.M., K.G.D., and D.G.B.; K.G.D. performed the computational analyses with input from D.G.B., V.E.M., M.P.V., and M.D.T.; D.G.B. and K.G.D. wrote the paper, with input from all other co-authors; K.G.D. and P.M.D. wrote the supplementary information, with input from all other authors. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** Raw reads and mitochondrial sequences have been deposited at the European Nucleotide Archive (ENA) with project number: PRJEB26011. Mitochondrial phylogenies are available at <https://osf.io/g5c8k/>.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/361/6397/85/suppl/DC1
Materials and Methods
Figs. S1 to S20
Tables S1 to S32
References (26–187)

11 January 2018; resubmitted 13 February 2018
Accepted 4 June 2018
10.1126/science.aas9411

Ancient goat genomes reveal mosaic domestication in the Fertile Crescent

Kevin G. Daly, Pierpaolo Maisano Delser, Victoria E. Mullin, Amelie Scheu, Valeria Mattiangeli, Matthew D. Teasdale, Andrew J. Hare, Joachim Burger, Marta Pereira Verdugo, Matthew J. Collins, Ron Kehati, Cevdet Merih Erek, Guy Bar-Oz, François Pompanon, Tristan Cumer, Canan Çakırlar, Azadeh Fatemeh Mohaseb, Delphine Decruyenaere, Hossein Davoudi, Özlem Çevik, Gary Rollefson, Jean-Denis Vigne, Roya Khazaeli, Homa Fathi, Sanaz Beizaei Doost, Roghayeh Rahimi Sorkhani, Ali Akbar Vahdati, Eberhard W. Sauer, Hossein Azizi Kharanaghi, Sepideh Maziar, Boris Gasparian, Ron Pinhasi, Louise Martin, David Orton, Benjamin S. Arbuckle, Norbert Benecke, Andrea Manica, Liora Kolska Horwitz, Marjan Mashkour and Daniel G. Bradley

Science **361** (6397), 85-88.
DOI: 10.1126/science.aas9411

How humans got their goats

Little is known regarding the location and mode of the early domestication of animals such as goats for husbandry. To investigate the history of the goat, Daly *et al.* sequenced mitochondrial and nuclear sequences from ancient specimens ranging from hundreds to thousands of years in age. Multiple wild populations contributed to the origin of modern goats during the Neolithic. Over time, one mitochondrial type spread and became dominant worldwide. However, at the whole-genome level, modern goat populations are a mix of goats from different sources and provide evidence for a multilocus process of domestication in the Near East. Furthermore, the patterns described support the idea of multiple dispersal routes out of the Fertile Crescent region by domesticated animals and their human counterparts.

Science, this issue p. 85

ARTICLE TOOLS

<http://science.sciencemag.org/content/361/6397/85>

SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2018/07/03/361.6397.85.DC1>

RELATED CONTENT

[file:/content](#)

REFERENCES

This article cites 122 articles, 14 of which you can access for free
<http://science.sciencemag.org/content/361/6397/85#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)